Lesson 11

Environmental Health Sciences Self-Study Course \$\$3010

Lesson 11: The Residential and Institutional Environment

Part I: Multiple Choice - APHA Evaluation

	The state of the s
1.	In the APHA method of measuring housing quality, which of the following would be considered a minor deficiency if found inadequate?
	a. central heating (b) water supply located outside of living unit c. lack of windows to the outside - basic d. minimum of two exits in case of fire (dual egress) - basic
2.	A minimum of of sleeping area per person is required in the APHA appraisal method.
	a. 10 ft ² b. 20 ft ² c. 40 ft ² d. 60 ft ² .
3.	In the APHA-PHS Recommended Housing Maintenance and Occupancy Ordinance, the responsibility of storing and disposing of all rubbish is assigned to the
	owner b. occupant c. landlord d. owner and occupant jointly.
4.	The ASHRAE recommended minimal thermal standard for dwellings is $\frac{24^{\circ}C}{16^{\circ}F}$ at 40% humidity and an air circulation rate of 45 fpm.
	a. 57°F b. 60°F c. 68°F d. 76°F.

Part I

Page 4

Fnvironmental	Haalth	Caiamana

Les	Son 11 Part I Page 5
	d. a backflow preventer properly vented.
	c) an open receptacle with an air gap
	b. a direct connection below floor level
	a. a grease trap
9.	The drain line of an ice machine should connect to the sewer system through
	d. 6 inches.
	c. 5 inches
(b) 4 inches
6	2 inches
	die diamage system.
	the drainage system.
8.	The waste discharge pipe from any appliance directly connected to the potable water system should terminate at least above the rim of a or receptacle directly connected to
	d. water meters.
1	c. air vents
,	7b, indirect waste piping
	a. gate valves
7.	Cross-connection controls include air gaps, backflow preventers, vacuum breakers and
	(d) developer.
	c. new owners
	b. local government
	a. local department of public works
6.	In unsewered areas, the responsibility for water and sewer for new subdivisions is assigned to the
	d. a dehumidifier in each unit.
	c. an air conditioner in each unit in tropical areas
	a) running water piped in b. 50 square feet of parking per unit
5.	Minimum APHA/PHS standards for housing include

	1.1		a aink should be
10. The air gap between the end of the f		rnow nm or un	e sink should be
times the diameter of the supply pig	e.		
a. 1 1/2			
6) 2			
c. 3			
d. 4.			
		- when the rela	tive humidity exceeds
11. Excessive condensation, corrosion,	and mildew occur	when the rela	ive limitalty exceeds
a. 20 percent			
b. 40 percent			
c. 80 percent			
60 percent.			
12. Vent gases combined with high mo	isture inside a chi	mney will form	n
12. Vent gases combined with high mo	isture miside a em		
a. a glass-like glaze			
b. hydrochloric acid			
c. sodium chloride			
d) sulfuric acid.			
(d.) suitaire acid.			
13. Nosocomial infections are a recurri	ng problem in		
13. Nosocomia infocuone are a session			
a. food establishments			
(b) hospitals			
c. swimming pools			
d. health spas.			
14. Minimum wash water temperature	in a hospital laun	dry is	
a. 146 to 150°F			
(b.) 160 to 167°F			
c. 170 to 178°F			
d. 185 to 196°F.			

- 15. It is recommended that a minimum of one water closet be provided for each _____ beds in a hospital or nursing home.
 - a. two
 - b. four
 - c. six
 - d eight.

Part II: Multiple Choice - Basic Housing

1. The housing code, regardless of who promulgates it, is (answer found in chapter 2)	
a complex environmental health code requiring the involvement of environmental health	h
 b. best left in building and housing departments not associated with the management of environmental health and protection issues c. not a good utilization of the expertise of the environmental health professional d. a program that does not require refined management at the 	
de l'emed management skills.	
 Lead levels exceeded 10ug/dL in U.S. children aged 6 and under in 1990. (answer found in MMWR March 12, 1993, Vol. 42/No. 9) 	
å. 300,000	
(b) 3,000	
c. 1,000,000 d. 3,000,000.	
 Utilization of will save the agency time and expense through the avoidance of extensive court time and to demonstrate that the agency uses legal action only as a measure of last resort. 	
(a) informal and formal hearings b. fines and tickets	
c. condemnation	
d. targeting low income areas for inspection.	
 In children, blood lead levels (BLLs) more than are associated with decreased intellectual performance and other adverse health events. 	
a. 2 ppm	
b. 5 μg/l	
® 10 μg/l	
d. 15 ppm.	

- 5. Lead poisoning is commonly associated with children living in homes built before
 - a. 1980
 - (b) 1970
 - c. 1950
 - d. 1940.
- 6. Most urban dwellers spend what percent of their total time indoors
 - a. 10 to 20 percent
 - b. 25 to 50 percent
 - © 80 to 90 percent
 - d. 70 to 80 percent.
- 7. Which of the following is the primary cause of indoor air quality problems?
 - (a) inadequate ventilation
 - b. contaminated outside air
 - c. radon
 - d. ragweed.
- CDC advocates a multitiered approach for environmental management of lead threats.
 This approach has which of the following steps
 - a. investigation and reduction of hazards
 - b. near universal screening
 - c. identification and remediation of hazards
 - (d.) all of the above.
- 9. Radon is detected in a home through the use of instrumentation that measures
 - a. geiger rays
 - beta particles
 - c. alpha particles
 - d. UV rays.

10. The major potential entry sources of radon are
a. cracks in concrete floors b. floor drains c. drainage sumps all of the above.
11. Radon can not be reduced with which of the following devices
 a humidifier b. electrostatic precipitators c. ionizers d. none of the above.
12. The best control of formaldehyde air pollution in the home is
 a. avoidance of the use of formaldehyde containing products b. the coating and sealing of materials containing formaldehyde c. increasing the amount of outside air d) all of the above.
13. Short term exposure to formaldehyde levels above can cause serious health or life threatening reactions
a. 50 ppm b. 20 ppm c. 10 ppm d. 0.5 ppm.
14. Exposure to formaldehyde can cause
 a. irritation to the respiratory passages b. contact dermatitis c. disturbed sleep d) all of the above.

15. If	radon is a problem in your area the followin	g locations wo	ould be	rooms of hi	gher risk;
a.	areas with showers and flood drains				
b.	basements	•		40	

c. washroomsd all of the above.

 Office space should provide at least ____ per min per occupant of clean outside air at minimum where smoking is not prohibited.

a. 5 ft³
b. 10 ft³
© 20 ft³
d. 30 ft.

17. To prevent chimney backdrafts, the chimney must be constructed ____ inches above the highest point of the roof at a minimum.

a. 4 inches
b. 8 inches
24 inches
24 inches
25 inches
36" above Play voets

18. The high moisture content of vent gases create


a acids which eat away at the mortar in the chimney
b. C₂H₂0
c. possible explosion hazard due to combustible gas

c. possible explosion hazard due to combustible gas all of the above.

19. The number of mobile home units in the United States

a. was 2.9 M in 1970
b. was 3.9 M in 1980
c. was 5.2 M in 1987
d. all of the above.

20. Most mobile homes	20.	Most	mobile	homes	are
-----------------------	-----	------	--------	-------	-----




- b. 33 percent are 65 years old or older
- c. 33 percent are 35 years old or younger
- (d) all of the above.

21. Mobile homes have been in general production since

- a. 1943
- b. 1954
- (2) 1965
- d. 1970.

22. The reasons for nosocomial infections in hospitals, and nursing homes are

- a. older patients with chronic diseases
- b. increased numbers of high risk patients
- c. invasive procedures
- (d) all of the above.
- 23. Which of the following are not typically members of the infection control committee
 - a hospital CEO
 - b. environmental control officer
 - c., infection control nurse



- 24. Which or the following are the most frequent disease transmission routes in child day care centers?
 - a. respiratory
 - b. fecal oral route
 - c. intimate contact
 - (d.) all of the above.

- 25. Individuals assigned to regulatory programs should be
 - a. exceptionally qualified by both training and experience
 - b. have degrees in institutional control
 - c. have achieved appropriate certifications
 - (a) a and c.

Part III: Biohazard, Questions Taken from Chapter 5, 1994 Supplement Multiple Choice	t,
1. A biohazard is	
 a. any material that negatively impacts a biological organism a material of biological composition, especially if infective, that constitutes a threat man or his environment c. a chemical that constitutes a threat to man or his environment d. any virus that causes fatal disease. 	at to

2.	The control of bloodborne pathogens is regulated by
	a. OSHA
	b. USDA

- c. NIH CDC.
- 3. Muerto Canyon virus (MCV) is a Hantavirus mainly carried by a. woodchucks
 - b. beavers (c.) deer mice
 - d. skunks.
- The Muerto Canyon virus causes disease of the _____ in humans.
 - (a) lungs b. kidneys
 - c. intestine
 - d. brain.

Envisor	
Environmental Health Sciences	
 The incidence of tuberculosis in English laborat was reported to be higher than for the 	OTV works
was reported to be higher than for the	general population.
a. 2 times	
b. 3 times	
c. 5 times	
(d) 10 times.	
6. Universal precautions refer to	
a. specific respiratory protection used in hospital	
an infection control measure in which all humans as infectious	an blood and
as infectious	in blood and certain body fluids are treated
c. procedures used when handling hazardous week	tes
d. none of the above.	
7. The biosafety containment level suitable for work hazard to personnel and the environment is	involving agents of moderate potential
hazard to personnel and the environment is	anderate potential
biosafety level 1 - orgs. that are biosafety level 2	not known to cause disease in healthy
c. biosafety level 3 - infectious agent while d. biosafety level 4 dangerous or exotic	ch may carete serious or potentially letal discuss agents that pose high risk of aerosol- itory infections + life-threatening discussions
The term containment is used in describing	tory infections + life-threatening fisces
a. control of infectious wastes in the laboratory b. a biosafety cabinet	
(c) safe methods for managing in f	
safe methods for managing infectious agents in the d. facilities designed for research on recombinant D	ne laboratory
Biological safety equipment includes	IVA.
a. biosafety cabinets - most common = Ch	iss I stype ABSC (BSL-2/3)
b. needles and syringes	
d a and c.	uns, eye protection true protection
Pipetting device	es, safety contrituge cups + rotors
101	

Environmental Health Sciences	
10. The four biosafety containment levels consist of a combination of	_
a. laboratory practices, safety equipment, and facilities laboratory techniques, research protocols, and safety equipment c. research protocols, safety training, and laboratory facilities d. laboratory practices, safety training, and research protocols.	
11. The protection of personnel and the immediate laboratory environment from exposure to infectious agents by good microbiological technique and safety equipment is called	
(a) primary containment b. secondary containment - fectives assoc, which surveyed the parties containment - none to prevent relyage in the event of d. none of the above.	r)
 Campylobacter (C. jejuni/C. coli, C. fetus subsp. fetus) may be safely used in the laboratory as a minimum at 	
a. biosafety level 1 biosafety level 2 c. biosafety level 3 d. biosafety level 4.	
13. The release of genetically engineered materials to the environment is regulated by	
a USDA and EPA b. NIH and CDC c. EPA and AEC d. FDA and PHS.	
14. The Medical Waste Tracking Act of 1988 required to establish a two year demonstration program for tracking medical waste in selected states	
a. CDC b. USDA/APHIS C EPA d. none of the above.	

15. Those laboratory activities that may result in personnel exposure to aerosols may result in	
a. CDC investigating b increased containment - (bissafety ()) c. the appointment of a biosafety officer d. none of the above.	
16. A class II biological safety cabinet (BSC) is designed to protect	
a. the work being done in the BSCb. the personnel using the BSC	

- 17. Biological safety cabinets should be performance tested every
 - a, 1 year

c. the environment all of the above.

- b 1 month
- c. 2 years
- d. 10 years.
- 18. A national performance standard has been developed by
 - a. NSF -
 - b. NSC
 - O NIH
 - d. ABSA.
- 19. The development and maintenance of an exposure control plan is one requirement of
 - a. EPA's biomedical waste regulations
 - OSHA's bloodborne pathogens regulations
 - c. USDA/APHIS plant genetics regulations
 - d. FIFRA.

- 20. A clean bench should be considered for use
 - a. as an alternative for a biosafety cabinet
 - b. as an alternative for a fume hood
 - only when there is no possible risk to personnel
 - d. when protecting the laboratory environment.
- 21. Perhaps the most important feature in implementing programs in biosafety is
 - a. the extensive use of rules and regulations
 - b. the reliance on outside contractors
 - c. working with the client in a consultative mode
 - none of the above.
- 22. The implementation of a biosafety program begins with
 - (a.) an assessment of risk
 - b. the development of a biosafety manual
 - c. training the laboratory personnel
 - d. consultation with CDC and NIH.
- 23. Herpesvirus simiae (B-virus) is naturally occurring in certain species of monkeys namely the macaque. The highest risk of humans acquiring this virus is from
 - a. primary cell cultures
 - (b) the bite of a macaque
 - c. aerosol exposure to saliva and urine
 - d. mouth pipetting in the laboratory.
- 24. Laboratories working with infectious agents have been shown to
 - (a) represent no threat to the community
 - b. represent a real and continuous threat to the community
 - c. represent a growing danger to the environment
 - d. none of the above.

- 25. The appointment and maintenance of an Institutional Biosafety Committee (IBC) is one of the requirements of
 - a. EPA
 - (b) CDC
 - c. NIH
 - d. FDA.